

Select country from dropdown menu

Ghana

GHA

Based on most recent available data.

Version 2.02.


2022

Output (coverage)

Outcome (service use/behaviour)

Impact

Condoms		Condom use with non-regular partners (%)		New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)	
Number of condoms distributed and sold / year (in millions)	id	Women 15-49	18	2010	18 000
Number of condoms distributed/sold per couple-year (age range 15-64)	id	Men 15-49	42	2015	16 000
Estimated condom distribution need met (%)	id			2020	14 000
Voluntary medical male circumcision (VMMC)		% of 2020 VMMC target achieved		HIV prevalence	
Number of VMMCs performed / year (in thousands)	na	Men	na	2020	4 400
% of annual VMMC target achieved	na	Men 15-24	96	2025	3 500
		Men 15-49	96		
ARV-based prevention		% of national PrEP need met			
Composite PrEP score (0-10)	3	All pop.	id		
Number of people who received PrEP at least once in the past 12 months	2135	Women 15+	id		
% of PLHIV on ART	71	Men 15+	id		
Key populations		% of PLHIV virally suppressed			
Sex workers (SW)		Condom use / last paid sex (%)			
Population size estimate for sex workers	id	Sex workers	id	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	89	Men 15-49	44	Sex workers all ages	6.9
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	99		
Gay men and other men who have sex with men (MSM)		Condom use / last anal sex (%)			
Population size estimate for men who have sex with men	54800	MSM	id	MSM <25 years	2.7
% of MSM who received at least two HIV prevention interventions (past 3 mo)	50	% on ART	95	MSM all ages	4.9
Prevention strategy includes core elements of MSM prevention package	> Half				
People who inject drugs (PWID)		% with safe injecting practices			
Population size estimate for people who inject drugs	id	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART	id		
Structural barriers and enablers		% of people who avoided health care because of stigma and discrimination			
Criminalization of sex work	Yes	Sex workers	id		
Criminalization of same-sex sexual acts	Yes	MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes	PWID	id		
Criminalization of transgender people	No	Transgender	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings		Condom use with non-regular partners (%)			
% of priority locations/districts with dedicated programs for AGYW & partners	id	Women 15-24	20	Women 15-24	1.1
Educational policies on HIV & sexuality education (secondary school)	Yes	Men 15-24	44	Men 15-24	0.4
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <14	% who completed lower secondary education	50	Adults, 15-49	1.7
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	id		
HIV testing services integrated with SRH services	Partial		id		



Summary Scores

AGYW	4
Sex workers	7
MSM	id
PWID	id
Condoms	id
VMMC	na
ART	7
PrEP	3

Acronyms: na ... not applicable; PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men; SRH ... sexual and reproductive health.

 Very good	 Low
 Good	 Very low
 Medium	 Insufficient data