

Select country from dropdown menu

Namibia

NAM

Based on most recent available data.

Version 2.02.

2022

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)		Outcome (service use/behaviour)		Impact		
Condoms						
Number of condoms distributed and sold / year (in millions)	0	Condom use with non-regular partners (%)	Women 15-49	66	<p>New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)</p> <p>2010: 9 500 2015: ~8 500 2020: 2 400 2025 target: 1 900</p>	
Number of condoms distributed/sold per couple-year (age range 15-64)	0		Men 15-49	82		
Estimated condom distribution need met (%)	0					
Voluntary medical male circumcision (VMMC)						
Number of VMMCs performed / year (in thousands)	16	% of 2020 VMMC target achieved	Men	45	<p>HIV prevalence</p> <p>2010: 14.0 2025: 29.9</p>	
% of annual VMMC target achieved	62	National male circumcision prevalence (%)	Men 15-24	22		
			Men 15-49	26		
ARV-based prevention						
Composite PrEP score (0-10)	3	% of national PrEP need met	All pop.	id	<p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>	
Number of people who received PrEP at least once in the past 12 months	18344	% of PLHIV virally suppressed	Women 15+	89		
% of PLHIV on ART	91		Men 15+	77		
Key populations						
<i>Sex workers (SW)</i>						
Population size estimate for sex workers	4000	Condom use / last paid sex (%)	Sex workers	42	Sex workers <25 years	14.0
% of SWs who received at least two HIV prevention interventions (past 3 mo)	id		Men 15-49	67	Sex workers all ages	29.9
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men	2200	Condom use / last anal sex (%)	MSM	55	MSM <25 years	3.6
% of MSM who received at least two HIV prevention interventions (past 3 mo)	88	% on ART	MSM LHIV	id	MSM all ages	7.8
Prevention strategy includes core elements of MSM prevention package	> Half					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs	id	% with safe injecting practices	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID LHIV	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	None	% on ART				
<i>Structural barriers and enablers</i>						
Criminalization of sex work	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same-sex sexual acts	Yes		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Criminalization of transgender people	No		Transgender	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	43	Condom use with non-regular partners (%)	Women 15-24	68	Women 15-24	6.2
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	84	Men 15-24	2.7
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <14	% who completed lower secondary education	Girls	62	Adults, 15-49	13.2
Provider-initiated condom promotion integrated into SRH services	No/id	% of women experienced physical or sexual violence from husband/partner	Women 15-19	id		
HIV testing services integrated with SRH services	Yes		Women 15-49	id		

Summary Scores

AGYW	5
Sex workers	id
MSM	7
PWID	id
Condoms	4
VMMC	5
ART	9
PrEP	3

Acronyms: na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

Very good
Good
Medium

Low
Very low
Insufficient data