

Select country from dropdown menu

Nigeria

NGA

Based on most recent available data.

Version 2.02.

2022

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)		Outcome (service use/behaviour)		Impact		
<b>Condoms</b>						
Number of condoms distributed and sold / year (in millions)	43	Condom use with non-regular partners (%)	Women 15-49	36		
Number of condoms distributed/sold per couple-year (age range 15-64)	1		Men 15-49	65		
Estimated condom distribution need met (%)	5					
<b>Voluntary medical male circumcision (VMMC)</b>						
Number of VMMCs performed / year (in thousands)	na	% of 2020 VMMC target achieved	Men	na		
% of annual VMMC target achieved	na	National male circumcision prevalence (%)	Men 15-24	96		
			Men 15-49	99		
<b>ARV-based prevention</b>						
Composite PrEP score (0-10)	8	% of national PrEP need met	All pop.	id		
Number of people who received PrEP at least once in the past 12 months	148952	% of PLHIV virally suppressed	Women 15+	93		
% of PLHIV on ART	90		Men 15+	89		
<b>Key populations</b>						
<i>Sex workers (SW)</i>						
Population size estimate for sex workers	874000	Condom use / last paid sex (%)	Sex workers	86	Sex workers <25 years	13.0
% of SWs who received at least two HIV prevention interventions (past 3 mo)	90		Men 15-49	74	Sex workers all ages	16.7
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	24		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men	240000	Condom use / last anal sex (%)	MSM	70	MSM <25 years	17.4
% of MSM who received at least two HIV prevention interventions (past 3 mo)	90	% on ART	MSM LHIV	26	MSM all ages	25.0
Prevention strategy includes core elements of MSM prevention package	> Half					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs	326100	% with safe injecting practices	PWID	36	PWID <25 years	11.8
% of PWID who received at least two HIV prevention interventions (past 3 mo)	76	% on opioid substitution therapy	PWID	id	PWID all ages	10.9
Prevention strategy includes core elements of PWID harm reduction package	> Half	% on ART	PWID LHIV	25		
<i>Structural barriers and enablers</i>						
Criminalization of sex work	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same-sex sexual acts	Yes		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Criminalization of transgender people	No		Transgender	id		
<b>Adolescent girls, young women (AGYW) &amp; partners in high-HIV incidence settings</b>						
% of priority locations/districts with dedicated programs for AGYW & partners	id	Condom use with non-regular partners (%)	Women 15-24	38	Women 15-24	0.7
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	62	Men 15-24	0.3
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <18	% who completed lower secondary education	Girls	59	Adults, 15-49	1.4
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19	13		
HIV testing services integrated with SRH services	Yes		Women 15-49	14		

Summary Scores

AGYW	5
Sex workers	8
MSM	8
PWID	2
Condoms	3
VMMC	na
ART	9
PrEP	8

Acronyms: na ... not applicable;  
 PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;  
 SRH ... sexual and reproductive health.

Very good	Low
Good	Very low
Medium	Insufficient data