


The State of HIV Prevention in Mozambique


2022

Number of new HIV infections (all ages)




2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.

Change in new HIV infections



HIV prevalence



Legend: Very good (Green), Good (Light Green), Medium (Yellow), Low (Orange), Very low (Red)

Policy and structural barriers

Key populations

Barrier	Sex workers	Gay men and other men who have sex with men	People who inject drugs	Transgender people	Prisoners
Criminalization of the behaviour of key populations	No	No	Yes	No	na
The national strategy includes critical elements of the programme package for key populations	> Half	> Half	Some	id	id
Avoided health care because of stigma and discrimination	id	id	id	id	na
Population size	224 000	64 000	14 000	id	id

Adolescent girls and young women

Indicator	15-19 years	15-49 years
Proportion of women who experienced intimate partner violence	id	id
Girls who completed lower secondary education	11%	
Policies on life skills-based HIV and sexuality education (secondary schools)	Yes	
Laws requiring parental consent for adolescents to access HIV testing services, age of consent	Yes, <12	

Linkages between HIV and sexual and reproductive health services

HIV testing services integrated within sexual and reproductive health	Yes
Provider-initiated condom promotion integrated into sexual and reproductive health services	Yes

HIV programme coverage and outcomes

ADOLESCENT GIRLS, YOUNG WOMEN & MALE PARTNERS

Condom use with a non-regular partner among young people 15-24 years old (%)

Group	Value
Young women	51
Young men	48

Target: 90%

Score: 31

KEY POPULATIONS

Sex workers

Condom use at last paid sex (%)

Group	Value
Sex workers	3
Clients	1

Target: 95%

Score: 36

Gay men and other men who have sex with men

Condom use at last anal sex (%)

Target: 90%

Score: 25

People who inject drugs

Use of harm reduction services (%)

Target: 40%

Score: 34

CONDOM PROGRAMMING

Condom use with a non-regular partner, 15-49 years (%)

Group	Value
Women	42
Men	47

Target: 90%

Score: 42

VOLUNTARY MEDICAL MALE CIRCUMCISION

Uptake of voluntary medical male circumcision

Group	Value
Women	44
Men	66

Target: 90%

Score: 20

ANTIRETROVIRAL DRUG-BASED PREVENTION

Pre-exposure prophylaxis

Number of people actively taking Pre-Exposure Prophylaxis (PrEP)

Year	Value
2017	600
2018	2000
2019	7434
2020	18513
2021	57717

Change in use of PrEP (2020-2021): +149%

Score: id

Antiretroviral treatment

People living with HIV virally suppressed (%)

Target: 73%

Score: id

PREVENTION OF VERTICAL TRANSMISSION OF HIV

MTCT rate

Score: id

% of HIV-positive pregnant women receiving ART

Score: id

Number of Estimated births to women living with HIV

Score: id

Number of new child infections due to vertical transmission

Score: id

Distribution of new child infections

- % of child infections because mother acquired HIV during pregnancy or breastfeeding: id
- % of child infections because mother did not receive antiretroviral therapy during pregnancy or breastfeeding: id
- % of child infections because mother did not continue antiretroviral treatment during pregnancy or breastfeeding: id
- % of child infections because mother was on antiretroviral treatment during pregnancy or breastfeeding, but was not virologically suppressed: id

Scores (1-10): Very good (Green), Good (Light Green), Medium (Yellow), Low (Orange), Very low (Red). id ... insufficient data, na ... not applicable

Data sources: UNAIDS 2021 epidemiological estimates; Global AIDS Monitoring 2021; and ICF - the DHS Program STATcompiler. Data sources for key population program coverage: Global AIDS Monitoring 2021, Global Fund and PEPFAR reports obtained in 2021. Note: The 2021 UNAIDS epidemiological estimates represent the year 2020. Other data points may refer to various years when the surveys were conducted. Note: 2020 and 2025 targets for reducing new HIV infections represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline. These reductions are required to achieve a 90% reduction by 2030.