

Select country from dropdown menu

South Africa

ZAF

Based on most recent available data.

Version 2.02.

2022

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)		Outcome (service use/behaviour)		Impact		
<b>Condoms</b>						
Number of condoms distributed and sold / year (in millions)	557	Condom use with non-regular partners (%)	Women 15-49	58	<p><b>New HIV infections</b> (ages 15+ trend vs. 2020 and 2025 targets)</p> <p>2010: 390 000 2015: ~300 000 2020: 210 000 2025 target: 77 000</p>	
Number of condoms distributed/sold per couple-year (age range 15-64)	28		Men 15-49	65		
Estimated condom distribution need met (%)	65					
<b>Voluntary medical male circumcision (VMMC)</b>						
Number of VMMCs performed / year (in thousands)	150	% of 2020 VMMC target achieved	Men	54	<p><b>HIV prevalence</b></p> <p>2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.</p>	
% of annual VMMC target achieved	19	National male circumcision prevalence (%)	Men 15-24	60		
			Men 15-49	id		
<b>ARV-based prevention</b>						
Composite PrEP score (0-10)	8	% of national PrEP need met	All pop.	id	<p><b>Summary Scores</b></p> <p>AGYW: 6</p> <p>Sex workers: 8</p> <p>MSM: 8</p> <p>PWID: 2</p> <p>Condoms: 6</p> <p>VMMC: 4</p> <p>ART: 7</p> <p>PrEP: 8</p>	
Number of people who received PrEP at least once in the past 12 months	346667	% of PLHIV virally suppressed	Women 15+	71		
% of PLHIV on ART	74		Men 15+	62		
<b>Key populations</b>						
<i>Sex workers (SW)</i>						
Population size estimate for sex workers	146000	Condom use / last paid sex (%)	Sex workers	id	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	69		Men 15-49	83	Sex workers all ages	45.0
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	70		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men	309700	Condom use / last anal sex (%)	MSM	72	MSM <25 years	id
% of MSM who received at least two HIV prevention interventions (past 3 mo)	79	% on ART	MSM LHIV	44	MSM all ages	42.9
Prevention strategy includes core elements of MSM prevention package	> Half					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs	82500	% with safe injecting practices	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	18	% on opioid substitution therapy	PWID	1	PWID all ages	19.5
Prevention strategy includes core elements of PWID harm reduction package	> Half	% on ART	PWID LHIV	41		
<i>Structural barriers and enablers</i>						
Criminalization of sex work	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same-sex sexual acts	No		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Criminalization of transgender people	No		Transgender	id		
<b>Adolescent girls, young women (AGYW) &amp; partners in high-HIV incidence settings</b>						
% of priority locations/districts with dedicated programs for AGYW & partners	50	Condom use with non-regular partners (%)	Women 15-24	61	Women 15-24	9.6
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	73	Men 15-24	3.1
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <12	% who completed lower secondary education	Girls	91	Adults, 15-49	17.0
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19	30		
HIV testing services integrated with SRH services	Yes		Women 15-49	30		

Acronyms: na ... not applicable;  
 PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;  
 SRH ... sexual and reproductive health.

Very good  
 Good  
 Medium  
 Low  
 Very low  
 Insufficient data