

Select country from dropdown menu

United Republic of Tanzania

TZA

Based on most recent available data.

Version 2.02.

2022

Output (coverage)

Outcome (service use/behaviour)

Impact

Output (coverage)		Outcome (service use/behaviour)		Impact	
Condoms					
Number of condoms distributed and sold / year (in millions)	72	Condom use with non-regular partners (%)	Women 15-49	28	New HIV infections (ages 15+ trend vs. 2020 and 2025 targets)
Number of condoms distributed/sold per couple-year (age range 15-64)	4		Men 15-49	35	
Estimated condom distribution need met (%)	17				
Voluntary medical male circumcision (VMMC)					
Number of VMMCs performed / year (in thousands)	571	% of 2020 VMMC target achieved	Men	100	HIV prevalence 2020 and 2025 targets represent the country's required contribution to global targets, a 75% reduction by 2020 and 82.5% reduction by 2025 against 2010 as a baseline.
% of annual VMMC target achieved	100	National male circumcision prevalence (%)	Men 15-24	82	
			Men 15-49	80	
ARV-based prevention					
Composite PrEP score (0-10)	6	% of national PrEP need met	All pop.	id	
Number of people who received PrEP at least once in the past 12 months	41335	% of PLHIV virally suppressed	Women 15+	91	
% of PLHIV on ART	86		Men 15+	75	
Key populations					
<i>Sex workers (SW)</i>					
Population size estimate for sex workers	id	Condom use / last paid sex (%)	Sex workers	72	Sex workers <25 years
% of SWs who received at least two HIV prevention interventions (past 3 mo)	20		Men 15-49	44	Sex workers all ages
Prevention strategy includes core elements of SW prevention package	> Half	% on ART	SWs LHIV	id	
<i>Gay men and other men who have sex with men (MSM)</i>					
Population size estimate for men who have sex with men	id	Condom use / last anal sex (%)	MSM	id	MSM <25 years
% of MSM who received at least two HIV prevention interventions (past 3 mo)	1	% on ART	MSM LHIV	id	MSM all ages
Prevention strategy includes core elements of MSM prevention package	> Half				
<i>People who inject drugs (PWID)</i>					
Population size estimate for people who inject drugs	36000	% with safe injecting practices	PWID	id	PWID <25 years
% of PWID who received at least two HIV prevention interventions (past 3 mo)	4	% on opioid substitution therapy	PWID LHIV	26	PWID all ages
Prevention strategy includes core elements of PWID harm reduction package	> Half	% on ART		id	
<i>Structural barriers and enablers</i>					
Criminalization of sex work	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	id	
Criminalization of same-sex sexual acts	Yes		MSM	id	
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id	
Criminalization of transgender people	No		Transgender	id	
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings					
% of priority locations/districts with dedicated programs for AGYW & partners	26	Condom use with non-regular partners (%)	Women 15-24	30	Women 15-24
Educational policies on HIV & sexuality education (secondary school)	Yes		Men 15-24	34	Men 15-24
Laws requiring parental consent for adolescents to access HIV testing services	Yes, <14	% who completed lower secondary education	Girls	27	Adults, 15-49
Provider-initiated condom promotion integrated into SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	Women 15-19	id	
HIV testing services integrated with SRH services	Yes		Women 15-49	id	

Summary Scores

AGYW	3
Sex workers	3
MSM	id
PWID	1
Condoms	2
VMMC	10
ART	9
PrEP	6

Acronyms: na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

SRH ... sexual and reproductive health.

Very good
Good
Medium

Low
Very low
Insufficient data